

Signaling Role of Hemocytes in *Drosophila* JAK/STAT-Dependent Response to Septic Injury

Hervé Agaisse,^{1,2,*} Ulla-Maja Petersen,^{1,3}
Michael Boutros,¹ Bernard Mathey-Prevot,³
and Norbert Perrimon^{1,2}

¹Department of Genetics
Harvard Medical School

²Howard Hughes Medical Institute
200 Longwood Avenue
Boston, Massachusetts 02115

³Dana-Farber Cancer Institute
Division of Pediatric Oncology and Hematology
44 Binney Street
Boston, Massachusetts 02115

Summary

To characterize the features of JAK/STAT signaling in *Drosophila* immune response, we have identified *totA* as a gene that is regulated by the JAK/STAT pathway in response to septic injury. We show that septic injury triggers the hemocyte-specific expression of *upd3*, a gene encoding a novel Upd-like cytokine that is necessary for the JAK/STAT-dependent activation of *totA* in the *Drosophila* counterpart of the mammalian liver, the fat body. In addition, we demonstrate that *totA* activation also requires the NF-KB-like Relish pathway, indicating that fat body cells integrate the activity of NF-KB and JAK/STAT signaling pathways upon immune response. This study reveals that, in addition to the pattern recognition receptor-mediated NF-KB-dependent immune response, *Drosophila* undergoes a complex systemic response that is mediated by the production of cytokines in blood cells, a process that is similar to the acute phase response in mammals.

Introduction

In mammals, local disturbances of physiological homeostasis such as trauma, injury, or bacterial infection lead to a systemic response known as the acute phase response (Baumann and Gauldie, 1994). A major early step in acute phase response is the local release of signaling molecules, such as cytokines, by activated macrophages, fibroblasts, and endothelial cells at the site of damage. Upon systemic release, these cytokines act distally from the site of damage and induce changes in behavior and biosynthetic capacities of a variety of target cells and organs in the whole organism. For instance, the liver is a central organ in acute phase response, and changes in hepatocyte gene expression profiles during acute phase response result in dramatic changes in serum concentrations of acute phase proteins, such as C-reactive protein (CRP) and serum amyloid A and P components (Steel and Whitehead, 1994). The cytokine IL6 has been identified as a major inducer of the metabolic changes that occur in liver during the acute phase

response (Fattori et al., 1994; Kopf et al., 1994). IL6 is produced at the site of damage in response to primary cytokines, such as IL1 or TNF (Sparacio et al., 1992; Zhang et al., 1990), or in response to the presence of bacterial cell wall compounds, such as lipopolysaccharide (LPS) (Zhang et al., 1994). Upon systemic release, IL-6 triggers JAK/STAT signaling in hepatocytes through activation of the cytokine class 1 receptor gp130 and the JAK kinases. This process ultimately leads to translocation of STAT dimers to the nucleus where they activate the transcription of target genes encoding acute phase proteins, such as C-reactive protein and LPS binding protein (LBP) (Alonzi et al., 2001; Schumann et al., 1996; Zhang et al., 1996).


In *Drosophila*, septic injury leads to the secretion of a battery of antimicrobial peptides by the fat body, an organ functionally equivalent to the liver in humans (Hoffmann et al., 1999). Two NF-KB-like signaling pathways are implicated in this process: the Toll and the Imd pathways (Lemaitre et al., 1995, 1996), which lead to the activation of the NF-KB-like transcriptional regulators Dif and Relish, respectively (Rutschmann et al., 2000; Stöven et al., 2000). While the Toll pathway is activated by gram-positive bacteria and fungi, activation of the Imd pathway is mediated by gram-negative bacteria (Lemaitre et al., 1997). Recently, the receptors involved in the specific recognition of gram-positive and gram-negative bacteria and leading to the activation of the Toll pathway and the Imd pathway, respectively, have been identified (Choe et al., 2002; Gottar et al., 2002; Michel et al., 2001; Ramet et al., 2002). These molecules belong to the peptidoglycan recognition protein (PGRP) family and are thought to mediate signaling in fat body by direct binding to the cell wall components of microorganisms upon infection.

While the mechanisms involved in NF-KB activation upon immune challenge are well documented both in insects and mammals, little is known about activation of the JAK/STAT pathway in insects. DNA binding activity and translocation of STAT in fat body cells in response to bacterial challenge has been documented in *Anopheles gambiae* (Barillas-Mury et al., 1999). Moreover, the immune challenge-inducible expression of the *Drosophila* thiolester-containing protein TEP1 has been shown to be constitutive in larvae displaying a gain-of-function mutation in JAK and reduced in a corresponding loss of function (Lagueux et al., 2000). These studies indicate that, as in mammals, the JAK/STAT pathway is activated upon immune challenge in *Drosophila*.


Here, we investigate the mechanisms involved in the JAK/STAT-dependent immune response to septic injury in *Drosophila*. First, we have identified *totA* as a gene that is regulated by the JAK/STAT pathway in fat body in response to septic injury. Then, we show that *totA* activation is mediated by the *Drosophila* homolog of the vertebrate cytokine class I receptor Domeless (Dome) and requires the hemocyte-specific expression of *upd3*, a gene encoding an Unpaired (Upd)-like cytokine. Finally, we show that *totA* activation also requires activation of the *relish* pathway in fat body. Altogether, our

*Correspondence: agaisse@rascal.med.harvard.edu

A



B



C




Figure 1. *totA* Expression Is JAK Kinase Dependent and Fat Body Specific

(A) Structure of the JAK/STAT pathway as required for embryonic development. The cytokine-like Upd binds the receptor Dome (Dome) resulting in receptor dimerization and activation of the receptor-associated JAK kinases Hopscotch (Hop) and tyrosine phosphorylation of the cytoplasmic domain of Dome. This creates docking sites for the SH2 domain of STAT. Upon recruitment to the active receptor complex, STATs become then phosphorylated by Hop and dimerize and translocate to the nucleus where STAT dimers activate transcription of target genes. The truncated version of Dome, Dome Δ CYT, lacks the intracellular region that is involved in signal transduction. This mutated receptor still contains the extracellular cytokine binding module and acts as a signaling antagonist, probably by titrating Upd.

(B) Transcriptional profiles of *drosomycin*, *diptericin*, and *totA* genes in wild-type and JAK mutant adult females. Northern blot analysis of total RNA extracted from adult flies 6 hr after infection with a mixture of gram-negative (*E. coli*) and gram-positive bacteria (*M. luteus*). Similar results were obtained for wild-type flies (*OreR*) grown at 24°C or 29°C. Flies displaying a JAK loss-of-function genotype (*M38/msv1*) were grown at 24°C. Flies displaying the thermosensitive JAK gain-of-function *Tum^L* were grown at 29°C.

(C) *totA* is expressed in fat body cells. *totA* expression was analyzed by in situ hybridization in abdominal adult fat body before (Control) and 6 hr after septic injury.

study indicates that *Drosophila* undergoes a complex pattern recognition-mediated and cytokine-mediated process in response to septic injury.

Results

totA Activation upon Septic Injury Relies on the JAK/STAT Pathway

In order to identify genes that are regulated by the JAK/STAT pathway in response to septic injury in adult flies, we screened for candidates that display an inducible expression upon immune challenge and that are constitutively expressed in flies carrying a gain-of-function mutation in the JAK/STAT pathway. To this end, we used custom-made cDNA microarrays (U.-M.P., unpublished data) to compare gene expression profiles of nonchallenged wild-type flies to gene expression profiles of challenged wild-type flies and to gene expression profiles of nonchallenged *Tum^L* flies displaying a gain-of-function mutation in the *Drosophila* JAK kinase Hopscotch (Figure 1A) (Harrison et al., 1995; Luo et al., 1995). We identified *MP1* as a gene that fulfilled both criteria for induction upon challenge and constitutive expression in a JAK/STAT gain-of-function mutation (Figure 1B). *MP1* expression was not induced in challenged flies displaying loss-of-function mutation in *hop* (*hop^{M38}/hop^{msv1}*

[Perrimon and Mahowald, 1986]), confirming the involvement of *Drosophila* JAK in *MP1* expression (Figure 1B). As control experiments, we also analyzed the expression of *drosomycin* and *diptericin*, which are controlled by the NF- κ B-like transcriptional regulators Dif and Relish, respectively. As expected, both loss-of-function and gain-of-function mutations in the JAK/STAT pathway did not affect *drosomycin* and *diptericin* expression in response to immune challenge, confirming the specific effects of JAK mutations on *MP1* expression (Figure 1B).

Sequence analysis of *MP1* cDNA revealed that *MP1* codes for TotA, a polypeptide that is produced by the larval fat body and accumulates in hemolymph in response to various stress conditions in flies (Ekengren et al., 2001; Ekengren and Hultmark, 2001). We next confirmed by in situ hybridization that *totA* expression was also mainly fat body specific in adult flies. As shown in Figure 1C, *totA* was weakly expressed in the fat body of unchallenged flies and strongly induced after septic injury.

STAT Is Activated in Fat Body Cells in Response to Septic Injury

Activation of the JAK/STAT pathway culminates in translocation of phosphorylated STAT dimers from the cytoplasm to the nucleus. In mosquitoes, it has been shown




Figure 2. Analysis of STAT Translocation in Fat Body Cells of Wild-Type and *JAK* Mutant Adult Flies

Flies were dissected 6 hr after infection with a mixture of gram-negative and gram-positive bacteria, and the fat body cells that remained attached to the abdominal cuticle were stained with the anti-DSTAT antibody (STAT protein, green) and propidium iodide (nuclear DNA, red). Similar results were obtained for wild-type flies (*OreR*) grown at 24°C or 29°C. Flies displaying a *JAK* loss-of-function genotype (*M38/msv1*) were grown at 24°C. Flies displaying the thermosensitive *JAK* gain-of-function *TumL* were grown at 29°C.

that AgSTAT, the homolog of *Drosophila* STAT, translocates into the nucleus of fat body cells in response to bacterial infection (Barillas-Mury et al., 1999). To further confirm the activation of the *JAK/STAT* pathway in *Drosophila* fat body in response to septic injury, we next examined the subcellular location of STAT protein in fat body cells by immunostaining. In unchallenged flies, STAT protein was located both in the cytoplasm and nucleus (Figure 2). In challenged flies, STAT had substantially cleared the cytoplasm and had accumulated in the nucleus (Figure 2). In contrast, there was no STAT nuclear translocation in *Drosophila JAK* mutant adult flies (*hop^{M38}/hop^{msv1}* [Perrimon and Mahowald, 1986]) (Figure 2). Conversely, a very strong staining was detected both in the cytoplasm and nucleus of flies carrying a *Drosophila JAK* gain-of-function mutation (*TumL* [Harrison et al., 1995; Luo et al., 1995]) (Figure 2). These results demonstrate that *totA* activation correlates with the *JAK*-dependent activation of STAT in *Drosophila* fat body cells in response to septic injury.

totA Expression Depends on *domeless* in Fat Body

Recently, a *Drosophila* homolog of the vertebrate cytokine class I receptor, Dome (a.k.a. Mom), has been identified (Brown et al., 2001; Chen et al., 2002). Mutations in *dome* result in embryonic defects similar to the embryonic phenotype associated with mutation in the *JAK/STAT* pathway components. A truncated version of Dome, Dome Δ CYT, has been generated by deletion of the intracellular region that is involved in signal transduction (Brown et al., 2001). This mutated receptor still contains the extracellular cytokine binding module and acts as a signaling antagonist, probably by titrating the ligand (Figure 1A). Accordingly, Dome Δ CYT overexpression during embryogenesis mimics the loss-of-function phenotype of *dome* mutants (Brown et al., 2001). To test whether Dome plays a role in *totA* expression, we used the GAL4/UAS system to express the dominant-negative form of Dome, Dome Δ CYT, in adult fat body. Northern blot analysis revealed that *totA* expression upon immune challenge was totally abolished in the corresponding animals (Figure 3A). We conclude that *totA* activation in response to bacterial infection is the result of a signaling event that is transduced by the *Drosophila* homolog of the vertebrate cytokine receptor, Dome, in the fat body.

The *upd/os* Locus Contains Three *upd*-like Genes

The involvement of *dome* in *totA* expression strongly suggested the existence of a cytokine-like molecule involved in the control of *totA* expression. *upd* has been characterized as a gene encoding the cytokine that activates the *JAK/STAT* pathway during *Drosophila* embryogenesis (Harrison et al., 1998). Strong alleles of *upd* are embryonic lethal, but weaker alleles, such as *outstretched* (*os*), give rise to adult flies that hold their wings at right angles and have small eyes. We found that *totA* activation was strongly decreased in *os* flies, suggesting that *upd* might be involved in *totA* expression (Figure 3B). However, *totA* activation was nearly wild-type in transheterozygous flies displaying the *os* mutation over a null mutation in *upd* (*upd^{YM55}*), suggesting that a defect in *upd* expression was not responsible for the lack of *totA* activation in the *os* genetic background (Figure 3B). Interestingly, *totA* activation was abolished in transheterozygous flies displaying the *os* mutation over a large deficiency (*os1A*) of the *upd* locus (Figure 3B). This suggests that the *os* mutation affects the expression of a gene involved in *totA* activation that maps to the *os/upd* locus, but that is not *upd*. Blast search analysis revealed the presence of two other *upd*-like cytokine-encoding genes at the *upd/os* locus (Figures 3C and 3D). *upd2* corresponds to CG5988 and maps 50 kb downstream from *upd* (Figure 3C). *upd3* corresponds to CG15062 (for the first and second exon) and CG5963 (for the third exon) and maps 25 kb downstream from *upd* (Figure 3C). These observations prompted us to hypothesize that *upd2* and/or *upd3* might be involved in *totA* expression (see below).

totA Expression in Fat Body Depends on *upd3* Expression in Hemocytes

To further investigate the potential role of *upd2* and/or *upd3* in *totA* expression, we first analyzed whether the




Figure 3. Involvement of *dome* and the *os* Locus in *totA* Expression
 (A) *totA* expression is *dome* dependent. Northern blot analysis of total RNA extracted from adult flies before challenge (0) or 6 hr after septic injury with a mixture of gram-negative and gram-positive bacteria. The effect of *Dome*ΔCYT overexpression was tested in flies displaying the UAS construct P(UAS-*dome*ΔCYT) and the enhancer trap P(C273-GAL4) driving GAL4 expression in fat body cells.
 (B) *totA* is not activated in *os* flies. Northern blot analysis of total RNA extracted from *OreR*, *os*, *os/updYM55* (a null *upd* allele), and *os/os1A* (a large deficiency covering the *os* locus) adult flies before (0) and 6 hr after septic injury.
 (C) Map of the *outstretched* (*os*) locus. *upd* corresponds to CG5993, *upd2* corresponds to CG5988, and *upd3* corresponds to CG15062 (for the first and second exons) and CG5963 (for the third exon). *os1A* is a deficiency covering the entire *os/upd* locus.
 (D) The Upd-like family members. Peptide sequence alignment of Upd, Om (1E) (from *Drosophila ananassae*), Upd2, and Upd3. SP indicates the signal peptide. Note that the N-terminal sequence of Upd3 is unknown. Asterisks indicate the position of introns. The four putative α helices are boxed.

expression of the *upd*-like genes was inducible upon septic injury. We failed to detect *upd2* expression in adult flies by using RT-PCR analysis. In contrast, we found that the level of *upd3* expression, which was very low in control animals, was significantly increased after septic injury (Figure 4A). We next analyzed the pattern of GFP expression in flies harboring a *upd3* promoter region-GAL4 fusion and a UAS-GFP reporter (see Experimental Procedures). We failed to detect any GFP production in fat body cells of control or challenged animals

(Figure 4B). However, we noticed that GFP production was strongly increased in blood cells after challenge, indicating that hemocytes might be the main site of *upd3* expression (Figure 4C). To investigate the functional importance of *upd3* hemocyte-specific expression in *totA* expression, we designed an in vivo RNAi strategy to silence *upd3* expression in a tissue-specific manner (Lee and Carthew, 2002). We used the *hemolymph-GAL4* and the *yolk-GAL4* constructs to drive the expression of the UAS-*iupd3* hairpin construct in hemocytes and fat body,




Figure 4. *upd3* Is Activated in Hemocytes in Response to Septic Injury

(A) RT-PCR analysis of total RNA extracted from *OreR* adult flies before challenge (0), 3, 6, and 9 hr after septic injury. RT-PCR signals were quantified using a PhosphorImager and normalized to the corresponding signal level of *rp49*. Results are expressed in arbitrary units, 100 corresponding to the level of *upd3* expression 3 hr after challenge.

(B and C) *upd3* expression is hemocyte specific. *Upd3* expression pattern was determined by analyzing GFP production (green) in fat body and hemocytes of *upd3-GAL4;UAS-GFP* flies before (Control) and after septic injury. Fat body tissue was visualized by DNA staining with propidium iodide (red), and hemocytes were identified by their ability to phagocytose fluorescent beads (red).

respectively (Figure 5A). While the production of *upd3* dsRNA in fat body did not interfere with *totA* expression, *upd3* dsRNA expression in hemocytes led to a strong decrease in *totA* activation upon septic injury (Figure

5B). Altogether, these experiments suggest that *upd3* activation in hemocytes subsequently leads to *totA* activation in the fat body.

totA Expression Is Regulated by the Relish Pathway in Fat Body

To further analyze the regulation of *totA* expression in fat body cells, we monitored *totA* expression in response to clean injury, septic injury with gram-negative bacteria (*E. coli*), or septic injury with gram-positive bacteria (*M. luteus*) (Figure 6A). Clean injury and septic injury with *M. luteus* resulted in a modest but significant induction of *totA* expression: 4-fold induction 6 hr after challenge and 7-fold induction 18 hr after challenge (Figure 6A). In sharp contrast, septic injury with *E. coli* resulted in a robust induction of *totA* expression: 25-fold induction at 6 hr and 35-fold induction at 18 hr (Figure 6A). Gram-negative bacteria therefore constitute the best inducer for *totA* expression. It is well established in flies that immune response to gram-negative bacteria is mediated by the Imd pathway through activation of TAK1 and the NF- κ B-like transcription factor Relish (Hedengren et al., 1999; Lemaitre et al., 1995; Vidal et al., 2001). We therefore analyzed *totA* expression in *TAK1* and in *relish* mutant flies. As shown in Figure 6B, *totA* activation after challenge was totally abolished in these mutants, indicating that, in addition to being JAK/STAT dependent, *totA* expression also requires the activity of the Relish pathway. We next analyzed whether the activity of the Relish pathway was specifically required in the fat body. To this end, we overexpressed *relish* dsRNA in fat body using the *UAS-irel* construct and the *yolk-GAL4* driver. As shown in Figure 6C, dsRNA-mediated silencing of *relish* expression leads to a failure in *totA* activation, indicating that Relish activity is specifically required in the fat body. Finally, we analyzed whether Relish activation in the fat body was sufficient to activate *totA* expression. It has been recently shown that overexpression of Imd in fat body cells leads to activation of Relish and therefore constitutive expression of the antimicrobial peptide genes, such as *dipthericin*, in absence of immune challenge (Georgel et al., 2001) (Figure 6C). As shown in Figure 6C, *totA* was not constitutively expressed in the corresponding flies, indicating that Relish activation is required in fat body but is not sufficient to activate *totA* expression.

Discussion

We have analyzed the activation of *totA*, a target of the JAK/STAT pathway in fat body, in response to septic injury. We demonstrate that septic injury triggers the activation of a cytokine-like molecule in blood cells, which in turn is necessary for activation of the JAK/STAT pathway in the fat body. Further, we show that the bacterial challenge-dependent activation of the NF- κ B pathway, which is necessary and sufficient to induce the expression of antibacterial peptides in the fat body, is also required for activation of *totA* in fat body cells. Altogether, our findings reveal the combined action of cytokine-mediated and pattern recognition-mediated processes in response to septic injury.





Figure 5. *upd3* Expression Is Required in Hemocytes for *totA* Fat Body Expression

(A) Hemolectin (*hml-GAL4*) and yolk (*yolk-GAL4*) drivers are hemocyte and fat body specific, respectively. Expression specificity was assessed by analyzing GFP production (green) in *hml-GAL4, UAS-GFP*, and *yolk-GAL4, UAS-GFP* flies. Fat body tissue was visualized by DNA staining with propidium iodide (red), and hemocytes were visualized by actin staining with phalloidin (red).

(B) Transcriptional profiles of *totA* and *diptericin* genes in flies displaying the *upd3*-silencing construct (*UAS-iupd3*). Northern blot analysis of total RNA extracted from *UAS-iupd3/+;TM3/+*, *UAS-iupd3/hml-GAL4*, and *UAS-iupd3/+;yolk-GAL4/+* flies before (0) and 6 hr after challenge.



TotA Is an Acute Phase Protein

To analyze the role of the JAK/STAT pathway during septic injury, we first identified a target gene of the pathway, *totA*. We showed that *totA* expression (1) takes place in the fat body, which is the insect equivalent of the mammalian liver; (2) is strongly induced in response to septic injury; and (3) requires the Dome/JAK/STAT signaling pathway in fat body and the expression of *upd3* in hemocytes. Similarly, the expression of the mammalian acute phase proteins (e.g., LBP and CRP) is modulated by activation of specific signaling pathways (e.g., the JAK/STAT pathway) in response to systemic release of cytokines (e.g., IL6) by activated macrophages (Schumann et al., 1996; Zhang et al., 1996). Altogether, our results demonstrate that TotA qualifies as a bona fide acute phase protein. In addition, *totC* and *totM* are also controlled by the JAK/STAT pathway upon septic injury, a characteristic that is probably shared by all the members of the *tot* family (H.A., unpublished data). Moreover, the *tot* family members are not the sole target of the JAK/STAT pathway upon septic injury. We have recently shown that expression of *CG11501*, a cys-

teine-rich polypeptide related to scorpion toxin, is also controlled by the JAK/STAT pathway (Boutros et al., 2002). These data indicate that the JAK/STAT pathway contributes to a global response upon immune challenge by controlling the expression of several acute phase proteins. As for most of their mammalian counterparts, the function of these *Drosophila* acute phase proteins is unclear. *totA* overexpression does not appear to protect NF-KB mutants, such as *key*, from gram-negative bacteria infection (H.A., unpublished data), indicating that TotA, unlike antimicrobial peptides such as Diptericin and Drosomycin, does not prevent bacterial growth. Altogether, these observations suggest that TotA is probably a general stress response factor involved in homeostasis of (damaged) tissues. Accordingly, it has been shown that TotA overexpression confers extended survival to flies subjected to heat stress, a treatment that certainly leads to disturbances of physiological homeostasis (Ekengren et al., 2001). Further in vivo characterization of acute phase protein function, such as TotA, using *Drosophila* as a model system will help us to understand the overall physiology of the acute phase response in insects and mammals.




Figure 6. *totA* Expression Relies on the *relish* Pathway in Fat Body
(A) *totA* is activated by septic injury with gram-negative bacteria. Northern blot analysis of total RNA extracted from *OreR* adult flies before challenge (Control), 6 hr, and 18 hr after clean injury (I), infection with *E. coli* (*Ec*), or infection with *M. luteus* (*MI*). Northern blot signals were quantified using a PhosphorImager and normalized to the mRNA level of ribosomal protein 49. Results are expressed in arbitrary units; 100 corresponds to the level of *totA* expression in *OreR* flies 18 hr after challenge with *E. coli* (*Ec*).
(B) *totA* expression relies on the *relish* pathway. Northern blot analysis of total RNA extracted from *OreR*, *TAK1*, or *relish* adult flies before challenge (0), 3, 6, and 9 hr after infection with *E. coli*.
(C) The *relish* pathway is required in fat body for *totA* expression. Northern blot analysis of total RNA extracted from *OreR* and flies displaying the fat body-specific driver *yolk-GAL4* and the *relish* silencing construct (*UAS-ire1*) before (0), 6, and 18 hr after infection with *E. coli*. The *UAS-imd* construct was used to activate the Relish pathway in fat body.

Upd3 and the Signaling Role of Hemocytes

Upd was first identified as a secreted molecule that activates the JAK/STAT pathway during *Drosophila* embryogenesis. Here, we provide evidence for the existence of a component of the JAK/STAT pathway: Upd3

that is produced in hemocytes in response to immune challenge. Although cytokine-like activities, such as IL1 and TNF α , have been previously reported as being produced by hemocytes from Lepidopteran larvae in response to LPS stimulation, none of these activities were shown to have a physiological function in vivo (Wittwer et al., 1999). To the best of our knowledge, *upd3* is the first example of a gene coding for a cytokine that is expressed in hemocytes and is required for signaling in fat body. This study therefore constitutes (to the best of our knowledge) the first demonstration that sentinel cells, such as hemocytes, play a signaling role in the *Drosophila* immune response. The nature of the signals that are detected by hemocytes and the signaling pathway(s) that trigger *upd3* activation in response to septic injury remain to be determined. Preliminary experiments indicated that *upd3* expression was severely impaired in *TAK1* flies after septic injury, suggesting that components of the Relish pathway (as defined in fat body cells) might be involved in *upd3* activation in hemocytes in response to bacterial infection (H.A., unpublished data). However, further analysis in *PGRP-LC* and *relish* mutant backgrounds were not consistent with this hypothesis. Clearly, the mechanisms involved in *upd3* regulation potentially constitute a new paradigm for studying the signals and the transduction machinery involved in the control of gene expression in activated hemocytes.

Integration of Signaling Pathway Activities in Fat Body

The NF-KB pathway is necessary and sufficient to activate the expression of antimicrobial peptides in response to septic injury in *Drosophila*. In contrast, *totA* activation in fat body requires both the JAK/STAT and the Relish pathways (Figure 7). Our study therefore reveals that fat body cells integrate various inputs, such as the presence of bacterial compounds (through NF-KB activation) and cytokines (through JAK/STAT activation), in response to septic injury (Figure 7). Interestingly, septic injury is not the only stimulus that activates *totA* expression. It has been shown that stress treatments, such as heat shock, dehydration, and mechanical pressure, also induce *totA* expression (Ekengren et al., 2001). Similarly, we have observed that injection of detergent solution into the hemolymph also triggers *totA* activation. Moreover, this response was strongly affected in both *hop* and *relish* mutants (H.A., unpublished data). Altogether, these observations indicate that *totA* regulation is achieved through detection of various stress stimuli and integration of the activity of at least two different signaling pathways, the JAK/STAT and the NF-KB pathways. We believe that such diversity in the mechanisms involved in the control of *totA* activation reflects the broad function of TotA as a general stress-response factor.

Concluding Remarks

In the past decade, *Drosophila* has emerged as a highly attractive model for the study of innate immunity. Remarkably, the mode of detection of microbial patterns through activation of pattern recognition receptors and NF-KB-like signaling pathways has been conserved throughout evolution. In addition to the detection of exogenous ligands such as LPS, the mammalian acute




Figure 7. Model for *totA* Activation in Response to Septic Injury
Septic injury triggers the activation of a cytokine-like molecule in hemocytes (Upd3), which in turn is necessary for activation of the JAK/STAT-dependent expression of *totA* in fat body. In addition, the gram-negative bacteria-triggered activation of the NF- κ B-like Relish pathway, which is necessary and sufficient to induce the expression of antibacterial peptide genes such as *dipterizin* (*dipt*), is also required for *totA* activation. *totA* expression thus results from the integration of pattern recognition receptor (PGRP-LC)- and cytokine (Upd3)-mediated responses in fat body cells.

phase response also involves the production of endogenous signaling molecules, such as cytokines. Our analysis indicates that a cytokine-mediated response also takes place in *Drosophila* in response to septic injury. This suggests that, very early in the evolution of multicellular organisms, a pattern recognition-mediated response was associated to a cytokine-mediated response. This association may reflect the need for integration at the organism level of a complex series of reactions that are executed by various organs not only to respond to exogenous stimuli, such as invading microorganisms, but also to orchestrate endogenous processes devoted to the maintenance of homeostasis.

Experimental Procedures

Fly Strains and Procedures

The *hop* alleles used (*M38*, *msv1*, and *TumL*) have been previously described (Harrison et al., 1995; Perrimon and Mahowald, 1986). *PGRP-LC* flies were provided by P. Manfruegli (Ramet et al., 2002), *TAK1* flies were provided by B. Lemaitre (Vidal et al., 2001), *relish* flies were provided by D. Hultmark (Hedengren et al., 1999), *P(UAS-dome Δ CYT)* line was obtained from J. Castelli-Gair Hombria (Brown et al., 2001), *P(UAS-*imd*)* and the *P(yolk-GAL4)* lines were provided by S. Naitza and P. Georgel (Georgel et al., 2001). The *P(hml-GAL4)* line used for hemocyte-specific expression has been generated by A. Goto (Goto et al., 2001) and is available from the Bloomington Center.

Infection Experiments

We used 1-week-old adult flies that were put on fresh medium and kept at room temperature 24 hr prior to challenge. Parameters such as temperature, humidity, food quality, population density, and age introduced considerable variability in *totA* expression. Septic injury was performed by pricking flies in the thorax using a thin needle previously dipped in a 10 \times concentrated overnight culture of *E. coli* and/or *M. luteus*.

Immunostainings

The 2.2 kb DNA fragment encoding the DSTAT protein was PCR amplified using a 4.0 kb cDNA clone (Hou et al., 1996) as a template and the following oligonucleotides as primers: GAAGAATTCATATGAGCTTGTGGAAGCGCATC, GGAGGATCCTCAAAGTTCTCAAAGTTTGT. The fragment was then ligated to the pBSK after digestion with EcoRI and BamHI. The fragment was then digested with NdeI and BamHI and subcloned into the NdeI and BamHI restriction sites of pET16b (Novagen), giving pET-STAT. *E. coli* strain BL21 was used to overexpress the DSTAT protein, using the T7 expression system. DSTAT protein was purified as inclusion bodies and injected into rat to generate anti-DSTAT antibodies. For immunostainings, flies were dissected in PBS 0.1% tween 4% formaldehyde, the remaining abdominal carcasses were incubated for 15 min in fresh PBS 0.1% tween 4% formaldehyde, washed twice in PBS 0.1% tween, and incubated overnight with the primary anti-DSTAT antibody PBS 0.1% tween 2% BSA at 4 $^{\circ}$ C. The samples were then washed three times in PBS 0.1% tween and incubated overnight with a FITC-conjugated anti-rat antibody (Jackson Lab.) in PBS 0.1% tween 2% BSA at 4 $^{\circ}$ C. The samples were then treated with RNase A (400 μ g/ml) and propidium iodide (20 μ g/ml) for 30 min, washed three times, and mounted in 80% glycerol antifade buffer (Molecular Probes). Fluorescent images were captured with a confocal microscope (Leica, TCS-NT) and subsequently processed using NIH image and Adobe Photoshop software.

RNA Preparation and Northern Blot Analysis

Adult flies were collected in eppendorf tubes, flash frozen in liquid nitrogen, and kept at -80° C. Total RNA extraction was performed with the RNA Trizol (GIBCO, BRL) method. Northern blot analysis was performed as previously described (Agaisse and Lereclus, 1996). The probes were generated by PCR amplification using genomic DNA as template and the following oligonucleotides as primers: *dipterizin*, GGAATCGGGGATTCTTTT, GACCAGAAAAGGAATCCC CGAT; *drosomycin*, ATCAAGTACTTGTTCGCCCTCTTCGCTGTC, ATATGTCGTAAGTAGTGGAGAGCTAAACGC; *rp49*, TCCGCCAG CATAAGGCCAGATCGT, TTAAGTCTTCTTGGAGAACGCAGCGC; and *totA* (*MP1*): TCATTCCGGTTTGTCTCAGCGTTCAAAAGTCA TAACC, CTCAAATATTAACAATATTAACCAGTGAATAATTGAG. Double-stranded DNA was labeled using the Readiprime II labeling kit (Amersham/Pharmacia).

In Situ Analysis of *totA* Expression

For in situ hybridization, flies were dissected in cold PBS, and the remaining abdominal carcasses were incubated overnight in PBS 8% formaldehyde at 4 $^{\circ}$ C. Hybridization was then performed as previously described (Hauptmann, 2001) using a RNA DIG-labeled probe generated by in vitro transcription, using the T7 polymerase and a pBS vector harboring the *totA* coding region (as described in Northern blot analysis section) as template.

RT-PCR Analysis

Twenty micrograms of total RNA were subjected to DNase I (Roche) treatment in 20 μ l DEPC-treated water for 30 min at 37 $^{\circ}$ C. Samples were then heated at 70 $^{\circ}$ C for 20 min. After addition of oligo dT (0.5 ng/ μ l) (Invitrogen), samples were heated at 70 $^{\circ}$ C for 5 min and quick cooled on ice. After addition of 10 μ l 5 \times RT-PCR buffer (Invitrogen), 4 μ l dNTP (10 mM), 5 μ l DTT, and 0.5 μ l RNasin (Roche), samples were incubated for 5 min at 37 $^{\circ}$ C. After addition of 2 μ l of Superscript Reverse Transcriptase (Invitrogen), samples were incubated for 2 hr at 42 $^{\circ}$ C. Samples were then diluted 100-fold, and 1 μ l of the dilution was subjected to PCR amplification using the following primers: *upd3*: ATGTCCAGTTTGCCTCT, TCGCTTGACAGAC TCTTA; and *rp49*: TCCGCCAGCATACAGGCCAGATCGT, TTAAGTCTTCTTGGAGAACGCAGCGC.

CGTTCTCTTGAGAACGCAGGCG. PCR cycles were performed as follows: 5 min at 94°C, then 30 s at 94°C, 30 s at 55°C, 1 min at 72°C for 20 cycles and, finally, 10 min at 72°C. Twenty microliters of a 100 µl PCR reaction were fractionated on a 1% agarose gel and then subjected to Southern blot analysis. In a typical experiment, as described in Figure 4, exposure time was 30 min at room temperature for *rp49* and 24 hr at -70°C for *upd3*. Signals were quantified by PhosphorImager and normalized to the corresponding signals for *rp49* expression. Results are expressed in arbitrary units; 100 corresponds to the maximum level of expression obtained for a given time course experiment.

Cloning and Transgene Constructs

The *upd3* promoter region was PCR amplified using the following oligonucleotides as primers: GGTGGTACCTCGTACAATGGTTTAA AAATAGCTCGCCAAAT and GGAGGATCCAGTGACCAGTTCCTG TTCAGGCGTCTCGTCGAT. The corresponding 4.0 kb DNA fragment was digested with KpnI and BamHI and cloned into the KpnI/BamHI restriction sites of pGATB (Brand and Perrimon, 1993). A 7.5 kb DNA fragment containing the *upd3* promoter region fused to the *GAL4* gene, and the *hsp70* polyadenylation signal sequence was generated by digestion with KpnI and NotI and subcloned into the KpnI and NotI sites of pCaSpeR4, giving pCaSpeR4: *upd3*'-*GAL4*.

The silencing constructs were generated by cloning sequentially into the NheI and AvrI restriction sites of pWIZ (Lee and Carthew, 2002) a 500 bp DNA fragment that was PCR amplified using specific oligonucleotides (displaying XbaI restriction site tails) as primers and an RT reaction of total RNA from challenged adults as template. Specific oligonucleotides were as follows: UAS-*iupd3*: GCTCTA GATCGCCTTGACAGACTCTTACGCTGGTTCG; GCTCTAGAATCGC GACCTGCAGATTTACGTGGCGCCT. UAS-*irel*: GCTCTAGACCGAC TTGCGGTTATTGATT; GCTCTAGATGGAACACATGGATCGCTAA.

Acknowledgments

We thank J. Castelli-Gair Hombria, D. Ferrandon, P. Georgel, D. Hultmark, P. Manfrueli, and S. Naitza for fly strains; and B. Lemaitre for fly and bacterial strains. We thank B. Lemaitre for helpful discussions and critical reading of the manuscript. We are grateful to J. Kopinja and C. Villalta for embryo injection. H.A. is a Research Associate at the Institut National de la Recherche Agronomique. This work was supported in part by a National Institutes of Health grant R01 HL62434 (to B.M.-P.). N.P. is an investigator of the Howard Hughes Medical Institute.

Received: February 27, 2002

Revised: July 9, 2003

Accepted: July 9, 2003

Published: September 8, 2003

References

Agaisse, H., and Lereclus, D. (1996). STAB-SD: a Shine-Dalgarno sequence in the 5' untranslated region is a determinant of mRNA stability. *Mol. Microbiol.* **20**, 633-643.

Alonzi, T., Maritano, D., Gorgoni, B., Rizzuto, G., Libert, C., and Poli, V. (2001). Essential role of STAT3 in the control of the acute-phase response as revealed by inducible gene inactivation [correction of activation] in the liver. *Mol. Cell.* **21**, 1621-1632.

Barillas-Mury, C., Han, Y.S., Seeley, D., and Kafatos, F.C. (1999). Anopheles gambiae Ag-STAT, a new insect member of the STAT family, is activated in response to bacterial infection. *EMBO J.* **18**, 959-967.

Baumann, H., and Gauldie, J. (1994). The acute phase response. *Immunol. Today* **15**, 74-80.

Boutros, M., Agaisse, H., and Perrimon, N. (2002). Sequential activation of signaling pathway in response to immune challenge in *Drosophila*. *Dev. Cell* **3**, 711-722.

Brand, A.H., and Perrimon, N. (1993). Targeted gene expression as a means of altering cell fates and generating dominant phenotypes. *Development* **118**, 401-415.

Brown, S., Hu, N., and Hombria, J.C. (2001). Identification of the first

invertebrate interleukin JAK/STAT receptor, the *Drosophila* gene domeless. *Curr. Biol.* **11**, 1700-1705.

Chen, H.W., Chen, X., Oh, S.W., Marinissen, M.J., Gutkind, J.S., and Hou, S.X. (2002). mom identifies a receptor for the *Drosophila* JAK/STAT signal transduction pathway and encodes a protein distantly related to the mammalian cytokine receptor family. *Genes Dev.* **16**, 388-398.

Choe, K.M., Werner, T., Stoven, S., Hultmark, D., and Anderson, K.V. (2002). Requirement for a peptidoglycan recognition protein (PGRP) in Relish activation and antibacterial immune responses in *Drosophila*. *Science* **296**, 359-362.

Ekengren, S., and Hultmark, D. (2001). A family of Turandot-related genes in the humoral stress response of *Drosophila*. *Biochem. Biophys. Res. Commun.* **284**, 998-1003.

Ekengren, S., Tryselius, Y., Dushay, M.S., Liu, G., Steiner, H., and Hultmark, D. (2001). A humoral stress response in *Drosophila*. *Curr. Biol.* **11**, 1479.

Fattori, E., Cappelletti, M., Costa, P., Sellitto, C., Cantoni, L., Carelli, M., Faggioni, R., Fantuzzi, G., Ghezzi, P., and Poli, V. (1994). Defective inflammatory response in interleukin 6-deficient mice. *J. Exp. Med.* **180**, 1243-1250.

Georgel, P., Naitza, S., Kappler, C., Ferrandon, D., Zachary, D., Swimmer, C., Kopczynski, C., Duyk, G., Reichhart, J.M., and Hoffmann, J.A. (2001). *Drosophila* immune deficiency (IMD) is a death domain protein that activates antibacterial defense and can promote apoptosis. *Dev. Cell* **1**, 503-514.

Goto, A., Kumagai, T., Kumagai, C., Hirose, J., Narita, H., Mori, H., Kadowaki, T., Beck, K., and Kitagawa, Y. (2001). A *Drosophila* haemocyte-specific protein, hemolectin, similar to human von Willebrand factor. *Biochem. J.* **359**, 99-108.

Gottar, M., Gobert, V., Michel, T., Belvin, M., Duyk, G., Hoffmann, J.A., Ferrandon, D., and Royet, J. (2002). The *Drosophila* immune response against Gram-negative bacteria is mediated by a peptidoglycan recognition protein. *Nature* **416**, 640-644.

Harrison, D.A., Binari, R., Nahreini, T.S., Gilman, M., and Perrimon, N. (1995). Activation of a *Drosophila* Janus kinase (JAK) causes hematopoietic neoplasia and developmental defects. *EMBO J.* **14**, 2857-2865.

Harrison, D.A., McCoon, P.E., Binari, R., Gilman, M., and Perrimon, N. (1998). *Drosophila* unpaired encodes a secreted protein that activates the JAK signaling pathway. *Genes Dev.* **12**, 3252-3263.

Hauptmann, G. (2001). One-, two- and three-color whole-mount in situ hybridization to *Drosophila* embryos. *Methods* **23**, 359-372.

Hedengren, M., Asling, B., Dushay, M.S., Ando, I., Ekengren, S., Wihlborg, M., and Hultmark, D. (1999). Relish, a central factor in the control of humoral but not cellular immunity in *Drosophila*. *Mol. Cell* **4**, 827-837.

Hoffmann, J.A., Kafatos, F.C., Janeway, C.A., and Ezekowitz, R.A. (1999). Phylogenetic perspectives in innate immunity. *Science* **284**, 1313-1318.

Hou, X.S., Melnick, M.B., and Perrimon, N. (1996). Marelle acts downstream of the *Drosophila* HOP/JAK kinase and encodes a protein similar to the mammalian STATs. *Cell* **84**, 411-419.

Kopf, M., Baumann, H., Freer, G., Freudenberg, M., Lamers, M., Kishimoto, T., Zinkernagel, R., Bluethmann, H., and Kohler, G. (1994). Impaired immune and acute-phase responses in interleukin-6-deficient mice. *Nature* **368**, 339-342.

Lagueux, M., Perrodou, E., Levashina, E.A., Capovilla, M., and Hoffmann, J.A. (2000). Constitutive expression of a complement-like protein in toll and JAK gain-of-function mutants of *Drosophila*. *Proc. Natl. Acad. Sci. USA* **97**, 11427-11432.

Lee, Y.S., and Carthew, R.W. (2002). Making a better RNAi vector for *Drosophila*: Use of intron spacers. *Methods* **30**, 322-329.

Lemaitre, B., Kromer-Metzger, E., Michaut, L., Nicolas, E., Meister, M., Georgel, P., Reichhart, J.M., and Hoffmann, J.A. (1995). A recessive mutation, immune deficiency (*imd*), defines two distinct control pathways in the *Drosophila* host defense. *Proc. Natl. Acad. Sci. USA* **92**, 9465-9469.

Lemaitre, B., Nicolas, E., Michaut, L., Reichhart, J.M., and Hoffmann,

- J.A. (1996). The dorsoventral regulatory gene cassette *spatzle/Toll/cactus* controls the potent antifungal response in *Drosophila* adults. *Cell* 86, 973–983.
- Lemaitre, B., Reichhart, J.M., and Hoffmann, J.A. (1997). *Drosophila* host defense: differential induction of antimicrobial peptide genes after infection by various classes of microorganisms. *Proc. Natl. Acad. Sci. USA* 94, 14614–14619.
- Luo, H., Hanratty, W.P., and Dearolf, C.R. (1995). An amino acid substitution in the *Drosophila* *hopTum-I* Jak kinase causes leukemia-like hematopoietic defects. *EMBO J.* 14, 1412–1420.
- Michel, T., Reichhart, J.M., Hoffmann, J.A., and Royet, J. (2001). *Drosophila* Toll is activated by Gram-positive bacteria through a circulating peptidoglycan recognition protein. *Nature* 414, 756–759.
- Perrimon, N., and Mahowald, A.P. (1986). *I(1)hopscotch*, A larval-pupal zygotic lethal with a specific maternal effect on segmentation in *Drosophila*. *Dev. Biol.* 118, 28–41.
- Ramet, M., Manfrulli, P., Pearson, A., Mathey-Prevot, B., and Ezekowitz, R.A. (2002). Functional genomic analysis of phagocytosis and identification of a *Drosophila* receptor for *E. coli*. *Nature* 416, 644–648.
- Rutschmann, S., Jung, A.C., Hetru, C., Reichhart, J.M., Hoffmann, J.A., and Ferrandon, D. (2000). The Rel protein DIF mediates the antifungal but not the antibacterial host defense in *Drosophila*. *Immunity* 12, 569–580.
- Schumann, R.R., Kirschning, C.J., Unbehauen, A., Aberle, H.P., Knope, H.P., Lamping, N., Ulevitch, R.J., and Herrmann, F. (1996). The lipopolysaccharide-binding protein is a secretory class 1 acute-phase protein whose gene is transcriptionally activated by APRF/STAT3 and other cytokine-inducible nuclear proteins. *Mol. Cell. Biol.* 16, 3490–3503.
- Sparacio, S.M., Zhang, Y., Vilcek, J., and Benveniste, E.N. (1992). Cytokine regulation of interleukin-6 gene expression in astrocytes involves activation of an NF-kappa B-like nuclear protein. *J. Neuroimmunol.* 39, 231–242.
- Steel, D.M., and Whitehead, A.S. (1994). The major acute phase reactants: C-reactive protein, serum amyloid P component and serum amyloid A protein. *Immunol. Today* 15, 81–88.
- Stöven, S., Ando, I., Kadalayil, L., Engstrom, Y., and Hultmark, D. (2000). Activation of the *Drosophila* NF-kappaB factor Relish by rapid endoproteolytic cleavage. *EMBO Rep.* 1, 347–352.
- Vidal, S., Khush, R.S., Leulier, F., Tzou, P., Nakamura, M., and Lemaitre, B. (2001). Mutations in the *Drosophila* dTAK1 gene reveal a conserved function for MAPKKs in the control of rel/NF-kappaB-dependent innate immune responses. *Genes Dev.* 15, 1900–1912.
- Wittwer, D., Franchini, A., Ottaviani, E., and Wiesner, A. (1999). Presence of IL-1- and TNF-like molecules in *Galleria mellonella* (Lepidoptera) haemocytes and in an insect cell line *Fromestigmene acraea* (Lepidoptera). *Cytokine* 11, 637–642.
- Zhang, Y.H., Lin, J.X., and Vilcek, J. (1990). Interleukin-6 induction by tumor necrosis factor and interleukin-1 in human fibroblasts involves activation of a nuclear factor binding to a kappa B-like sequence. *Mol. Cell. Biol.* 10, 3818–3823.
- Zhang, Y., Broser, M., and Rom, W.N. (1994). Activation of the interleukin 6 gene by *Mycobacterium tuberculosis* or lipopolysaccharide is mediated by nuclear factors NF-IL6 and NF-kappa B. *Proc. Natl. Acad. Sci. USA* 91, 2225–2229.
- Zhang, D., Sun, M., Samols, D., and Kushner, I. (1996). STAT3 participates in transcriptional activation of the C-reactive protein gene by interleukin-6. *J. Biol. Chem.* 271, 9503–9509.